

Cross Sections for the Electron Activation of γ -Ray Fluorescence

Silviu Olariu* and Agata Olariu

National Institute of Physics and Nuclear Engineering, Tandem Laboratory
76900 Magurele, P.O. Box MG-6, Bucharest, Romania

Yoshiaki Ito and Takeshi Mukoyama

Institute for Chemical Research, Kyoto University, Uji, Kyoto 611, Japan

February 8, 2008

Abstract

We report cross sections for the direct excitation of γ -ray transitions up to 200 keV by the transient electromagnetic fields of electrons from a beam, for electron incident kinetic energies of 500 keV and 5 MeV. The cross sections for the electron activation of γ -ray fluorescence are of the order of 300 nb for an electron incident kinetic energy of 500 keV, and are of the order of 10 μ b for an electron incident kinetic energy of 5 MeV. The electron excitation of nuclear transitions may lead to the development of pulsed sources of gamma radiation of narrowly defined energy.

PACS numbers: 23.20.Lv, 23.20.Nx, 42.55.Vc

*e-mail: olariu@ifin.nipne.ro

It has been shown [1] that if a beam of incident electrons produces holes in the atomic shell of certain atomic species, this atomic excitation can be transferred to the nucleus. The nucleus thus raised in an excited state then decays by the emission of Mössbauer gamma-ray photons. A pulse of incident electrons would produce in this way a pulse of Mössbauer γ -ray photons, which have a narrowly-defined energy. The transfer of excitation from the atomic shell to the nucleus is due to the hyperfine interaction energy, and depends on the existence of a match between an atomic transition energy and a transition from the ground nuclear state to an excited nuclear state. The probability of the X-ray electron-nuclear transition (XENDT) is inversely proportional to the square of this detuning between the atomic transition energy and the nuclear transition energy.

In this work we study the *direct* excitation of γ -ray transitions up to 200 keV by the transient electromagnetic field of electrons from an incident beam. This type of nuclear excitation is no longer restricted by the requirement of the existence of a near resonance between the atomic and nuclear transition energies, and is in principle open to all nuclear species. Cross sections for the nuclear activation by electrons of γ -ray fluorescence have been calculated in this work using the results of Robl, [2] who gave expressions of the cross sections in the Born approximation, for magnetic and electric multipole transitions of arbitrary order. In these expressions, the nuclear properties are accounted for by the partial γ -ray widths of the excited Mössbauer state. The partial widths are calculated in this work from tabulated nuclear data on half-lives, relative intensities and internal conversion coefficients.

The cross sections for the electron activation of γ -ray fluorescence are of the order of 300 nb for incident electrons with a kinetic energy of 500 keV, being about two orders of magnitude lower than the cross sections for the excitation of Mössbauer transitions via electron-nuclear interactions, [1] but the number of nuclei which can be excited directly by electrons is larger, because of the absence of the near-resonance condition of [1]. The cross sections for the electron activation of γ -ray fluorescence are of the order of $10 \mu\text{b}$ for incident electrons with a kinetic energy of 5 MeV. The γ -ray activity induced by an electron beam incident on a sample containing the nuclear species under consideration is of the order of 10^4 Bq/mA, for an electron incident kinetic energy of 500 keV. Much larger activities could be produced if the incident electron beam consists of *pulses* of electrons. Such pulsed sources would produce γ -ray photons of narrowly defined energy. Because of their narrowly-defined energy, the γ -ray pulses produced in this way might be used for interference experiments, such as γ -ray holography. [3] Pulsed sources of Mössbauer radiation would be also valuable for the cases when the half-life of a conventional Mössbauer source is too short for practical use.

The cross section for the excitation by an incident electron of an EL transition from a ground nuclear state $|g\rangle$ to an excited nuclear state $|e\rangle$ can be written in the Born approximation as [2]

$$\sigma_{EL} = \frac{2I_e + 1}{2I_g + 1} \frac{\hbar \Gamma_{eg}^{EL}}{mc^2} F_{EL}, \quad (1)$$

where Γ_{eg}^{EL} is the partial width for the emission of a γ -ray photon of multipolarity EL in the transition $|e\rangle \rightarrow |g\rangle$, I_e, I_g are the spins of the excited and ground nuclear state, and m is the electron mass. The quantity F_{EL} is given by

$$\begin{aligned} F_{EL} = & \frac{1}{L+1} \frac{\pi \alpha \hbar^2 c^2 m}{E_\gamma^3 p_0^2} \left[\frac{1}{4} (L-1)(E_0 + E)^2 K_{L-2} \right. \\ & + \left(E_0 E - (L+1)m^2 c^4 - \frac{3}{2} L(E_0^2 + E^2) \right) K_{L-1} \\ & \left. + \left(2LE_0 E + \frac{1}{4}(7L+1)E_\gamma^2 \right) K_L - \frac{1}{2} L E_\gamma^2 K_{L+1} \right], \end{aligned} \quad (2)$$

where α is the fine structure constant, E_γ is the energy of the γ -ray photon emitted in the transition $|e\rangle \rightarrow |g\rangle$, E_0 is the relativistic energy of the incident electron including rest mass, p_0 is the momentum of the incident electron, $p_0 = (E_0^2 - m^2 c^4)^{1/2}/c$, and E is the energy of the electron after excitation of the nucleus, $E = E_0 - E_\gamma$. The dimensionless quantities K_L are defined by the recurrence relation

$$K_{L+2} - 2K_{L+1} + K_L = \frac{1}{L+1} (x_2^{L+1} - x_1^{L+1}), \quad (3)$$

with

$$K_0 = \frac{p_0 p}{m^2 c^2}, \quad K_1 = K_0 + 2 \ln \frac{mc^2 E_\gamma}{E_0 E - m^2 c^4 - p_0 p c^2}, \quad (4)$$

where p is the momentum of the electron after excitation of the nuclear state, $p = (E^2 - m^2 c^4)^{1/2}/c$, and the dimensionless quantities x_1, x_2 are

$$x_1 = \frac{(p_0 - p)^2}{\hbar^2 k^2}, \quad x_2 = \frac{(p_0 + p)^2}{\hbar^2 k^2}. \quad (5)$$

The cross section for the excitation by an incident electron of an ML transition from a ground nuclear state $|g\rangle$ to an excited nuclear state $|e\rangle$ can be written in the Born approximation as [2]

$$\sigma_{ML} = \frac{2I_e + 1}{2I_g + 1} \frac{\hbar \Gamma_{eg}^{ML}}{mc^2} F_{ML}, \quad (6)$$

where Γ_{eg}^{ML} is the partial width for the emission of a γ -ray photon of multipolarity ML in the transition $|e\rangle \rightarrow |g\rangle$, and the quantity F_{ML} is given by

$$F_{ML} = \frac{\pi \alpha \hbar^2 c^2 m}{E_\gamma^3 p_0^2} \left[(E_0 E - m^2 c^4) K_L + \frac{1}{4} E_\gamma^2 (K_{L+1} - x_1 x_2 K_{L-1}) \right]. \quad (7)$$

The partial γ -ray width Γ_{eg} of the transition $|e\rangle \rightarrow |g\rangle$ can be calculated as

$$\Gamma_{eg} = \frac{\ln 2}{t_e} \frac{R_{eg}}{\sum_l (1 + \alpha_{el} R_{el})}, \quad (8)$$

where t_e is the half-life of the excited state $|e\rangle$, the index l designates all nuclear states lower than $|e\rangle$, including the ground state $|g\rangle$, R_{el} is the relative intensity of the transition $|e\rangle \rightarrow |g\rangle$, and α_{el} are the internal conversion coefficients. Among the cases studied in this work, we encountered mixed transitions only of the $M1 + E2$ type. In the case of mixed $M1 + E2$ transitions, the internal conversion coefficients have been calculated as

$$\alpha = \frac{1}{1 + \delta^2} \alpha_{M1} + \frac{\delta^2}{1 + \delta^2} \alpha_{E2}, \quad (9)$$

the partial widths are

$$\Gamma_{eg}^{M1} = \frac{1}{1 + \delta^2} \Gamma_{eg}, \quad \Gamma_{eg}^{E2} = \frac{\delta^2}{1 + \delta^2} \Gamma_{eg}, \quad (10)$$

and the total cross section for the electron excitation of the $|e\rangle$ state is

$$\sigma_{M1+E2} = \sigma_{M1} + \sigma_{E2}, \quad (11)$$

where δ is the mixing ratio for the $M1$ and $E2$ components of the γ -ray transition $|e\rangle \rightarrow |g\rangle$. [5] A pure $M1$ transition corresponds to $\delta = 0$, and a pure $E2$ corresponds formally to $\delta = \infty$. For the $E1$, $M2$ and $E3$ transitions encountered in this work, we had $\Gamma_{eg}^{E1} = \Gamma_{eg}$, $\Gamma_{eg}^{M2} = \Gamma_{eg}$, and $\Gamma_{eg}^{E3} = \Gamma_{eg}$.

The cross section σ_γ for the emission of a γ -ray photon to be induced by an incident electron is lower than σ_{EL} , σ_{ML} or σ_{M1+E2} by the factor $f = \Gamma_{eg}/\Gamma_e$, where $\Gamma_e = \ln 2/t_e$,

$$\sigma_\gamma = f \sigma_e, \quad (12)$$

where σ_e denotes σ_{EL} , σ_{ML} or σ_{M1+E2} , according to the case. In the case of a Mössbauer experiment, the cross section σ_γ must be multiplied by the recoil-free fraction.

The main limitation of the previous expressions for the electron excitation cross section arises from the condition of validity of the Born approximation, which requires that $Z\alpha(E/pc) \ll 1$, where Z is the proton number of the nucleus. As discussed by Robl, [2] there is also an upper limitation of the photon energy E_γ of about 10 MeV which is not of concern for the present work, because we are studying transitions of much lower energy.

We have calculated cross sections for 144 cases of excitation of nuclear states having energies less than 200 keV, connected to the ground state by $E1$, $M1$, $E2$, $M2$ or $E3$ transitions, for which the half-lives t_e , the relative intensities R_{el} and the mixing ratios were known. [4] We have calculated the internal conversion coefficients by interpolation [6], [7] and have also used values calculated on-line at the National Nuclear Data Center, Brookhaven. [8] The

cross section σ_e and σ_γ are listed in Table I, for a kinetic energy of the incident electron of $E_0 - mc^2 = 500$ keV, and $E_0 - mc^2 = 5$ MeV. It can be seen from Table I that the largest cross section σ_γ for an electron kinetic energy of 500 keV for the emission of a γ -ray photon has the value 377 nb. The cross sections at this electron energy are about two orders of magnitude lower than the cross sections for the excitation of Mössbauer transitions via electron-nuclear interactions, [1] but the number of nuclear states which can be excited directly by electrons is larger, because of the absence of the near-resonance condition of [1]. It can be seen from Table I that the cross section for an incident kinetic energy of the electron of 5 MeV are significantly larger than the cross sections corresponding for a kinetic energy of 500 keV. We have also given in Table I values of the quantity $Z\alpha E/pc$.

As estimated in [1], when the incident electron beam is incident on a suitable target, such cross sections result in activities of the order of 10^4 Bq/mA for an electron incident kinetic energy of 500 keV. In the case of a *pulsed* incident electron beam the resulting activities would be much larger over the period of the pulse, because of the much larger incident current. The photons emitted by such a pulsed γ -ray source would have a narrowly defined energy, and would be suitable for interference experiments, such as holography. [3]

Electron activated γ -ray sources could also be of interest for Mössbauer experiments in cases where the half-life of conventional radioactive sources is too short to be practical. For example, among the cases listed in Table I, the radioactive Mössbauer sources which could be used for ^{162}Er , ^{188}Os , ^{164}Er , ^{165}Ho , ^{184}Os , ^{180}W , ^{187}Re , ^{180}Hf , ^{162}Dy , ^{174}Hf , ^{176}Yb , ^{187}Os , ^{160}Gd , ^{164}Dy , ^{163}Dy have a half-life less than 1 day, and no source is available for ^{40}K .

Although the cross sections of X-ray electron-nuclear double transitions discussed in [1] and the cross sections for the direct excitation of nuclear transitions are not very large, these processes appear to be of real interest for the development of electron-activated sources of γ -ray photons of narrowly defined energy.

Acknowledgment One of the authors (S.O.) acknowledges the financial support of the University of Kyoto during the preparation of this work.

References

- [1] S. Olariu, A. Olariu and V. Zoran, Phys. Rev. C **56**, 381 (1997).
- [2] H. R. Robl, Nucl. Phys. **2**, 641 (1956/57).
- [3] B. Adams, D. V. Novikov, T. Hiort and G. Materlik, Phys. Rev. B **57**, 7526 (1998).
- [4] *Table of Isotopes*, 8th ed., edited by R. Firestone (Wiley, New York, 1996).
- [5] Ref. [4], vol. I, p. XX.
- [6] I. M. Band, M. B. Trzhaskovskaya and M. A. Listengarten, Atomic Data and Nuclear Data Tables **18**, 433 (1976).
- [7] F. Rössel, H. M. Fries, K. Alder and H. C. Pauli, Atomic Data and Nuclear Data Tables **21**, 91 (1978), **21**, 291 (1978).
- [8] National Nuclear Data Center, Brookhaven, <http://www.nndc.bnl.gov>.

TABLE CAPTION

TABLE I. Cross section σ_e for the electron excitation of nuclear states and cross section σ_γ for the generation of fluorescence γ -ray photons of energy E_γ by electrons with incident kinetic energy $E_0 - mc^2 = 500$ keV, and $E_0 - mc^2 = 5$ MeV. The powers of 10 are represented with the aid of the symbol E, so that the value 3×10^{-10} is written as 3E-10.

nucleus	nat.ab. %	E_γ , keV	t_e , s	multipole $ e\rangle \rightarrow g\rangle$	δ	f	σ_e , nb 500 keV	σ_γ , nb 500 keV	$Z\alpha E/pc$, 500 keV	σ_e , nb 5 MeV	σ_γ , nb 5 MeV	$Z\alpha E/pc$, 5 MeV
¹⁶⁹ Tm	100.00	130.5	3E-10	E2		3.19E-1	1.18E3	3.77E2	0.618	3.54E4	1.13E4	0.506
¹⁵⁶ Dy	0.06	137.8	8.2E-10	E2		5.39E-1	6.87E2	3.7E2	0.594	2.09E4	1.12E4	0.484
¹⁵⁴ Gd	2.18	123.1	1.2E-9	E2		4.55E-1	7.33E2	3.33E2	0.571	2.17E4	9.85E3	0.469
¹⁶² Er	0.14	102.0	1.2E-9	E2		2.65E-1	1.16E3	3.08E2	0.600	3.3E4	8.75E3	0.498
¹⁵² Sm	26.70	121.8	1.4E-9	E2		4.61E-1	6.52E2	3.01E2	0.553	1.92E4	8.87E3	0.454
¹⁵⁰ Nd	5.64	130.2	1.4E-9	E2		5.36E-1	5.29E2	2.83E2	0.538	1.58E4	8.49E3	0.440
¹⁹⁰ Os	26.40	186.7	3.6E-10	E2		7.01E-1	3.98E2	2.79E2	0.707	1.33E4	9.32E3	0.557
¹⁸⁶ Os	1.58	137.2	8.2E-10	E2		4.35E-1	5.74E2	2.5E2	0.684	1.74E4	7.57E3	0.557
¹⁵⁸ Dy	0.10	98.9	1.7E-9	E2		2.59E-1	9.31E2	2.41E2	0.581	2.64E4	6.83E3	0.484
¹⁸⁶ W	28.60	122.6	1E-9	E2		3.58E-1	6.74E2	2.41E2	0.660	1.99E4	7.12E3	0.542
¹⁸⁸ Os	13.30	155.0	7.1E-10	E2		5.49E-1	4.34E2	2.38E2	0.691	1.36E4	7.46E3	0.557
¹⁶⁴ Er	1.61	91.4	1.5E-9	E2		1.91E-1	1.18E3	2.26E2	0.597	3.29E4	6.3E3	0.498
¹⁶⁵ Ho	100.00	94.7	2.2E-11	M1+E2	0.155	2.4E-1	8.53E2	2.05E2	0.589	1.44E4	3.46E3	0.491
¹⁸⁴ W	30.67	111.2	1.3E-9	E2		2.76E-1	7.18E2	1.98E2	0.656	2.08E4	5.73E3	0.542
¹⁵⁶ Gd	20.47	89.0	2.2E-9	E2		2.03E-1	9.55E2	1.94E2	0.561	2.66E4	5.38E3	0.469
¹⁸⁵ Re	37.40	125.4	1E-11	M1+E2	0.18	2.63E-1	7.21E2	1.9E2	0.670	1.2E4	3.17E3	0.550
¹⁸⁴ Os	0.02	119.8	1.2E-9	E2		3.16E-1	5.88E2	1.86E2	0.677	1.73E4	5.46E3	0.557
¹⁸⁰ W	0.13	103.6	1.3E-9	E2		2.24E-1	8.26E2	1.85E2	0.653	2.36E4	5.28E3	0.542
¹⁶⁸ Yb	0.13	87.7	1.5E-9	E2		1.55E-1	1.18E3	1.82E2	0.613	3.27E4	5.06E3	0.513
¹⁶⁰ Dy	2.34	86.8	2E-9	E2		1.76E-1	1.03E3	1.8E2	0.578	2.85E4	5E3	0.484
¹⁸² W	26.30	100.1	1.4E-9	E2		2.02E-1	8.31E2	1.68E2	0.652	2.36E4	4.76E3	0.542
¹⁸⁷ Re	62.60	134.2	1.1E-11	M1+E2	0.175	3.03E-1	5.53E2	1.67E2	0.674	8.74E3	2.64E3	0.550
¹⁷⁸ Hf	27.30	93.2	1.5E-9	E2		1.74E-1	9.6E2	1.67E2	0.633	2.69E4	4.67E3	0.528
¹⁸⁰ Hf	35.10	93.3	1.5E-9	E2		1.75E-1	9.44E2	1.65E2	0.633	2.65E4	4.62E3	0.528
¹⁸¹ Ta	99.99	136.3	3.9E-11	M1+E2	0.41	3.6E-1	4.41E2	1.59E2	0.656	1.13E4	4.06E3	0.535
¹⁷⁶ Hf	5.21	88.3	1.4E-9	E2		1.45E-1	1.09E3	1.58E2	0.631	3.04E4	4.4E3	0.528
¹⁷⁰ Yb	3.05	84.3	1.6E-9	E2		1.35E-1	1.16E3	1.57E2	0.612	3.21E4	4.33E3	0.513
¹⁶⁶ Er	33.60	80.6	1.8E-9	E2		1.27E-1	1.21E3	1.53E2	0.594	3.31E4	4.19E3	0.498
¹⁶² Dy	25.50	80.7	2.2E-9	E2		1.38E-1	1.09E3	1.5E2	0.576	2.98E4	4.12E3	0.484
¹⁵⁸ Gd	24.84	79.5	2.5E-9	E2		1.43E-1	1.05E3	1.5E2	0.559	2.88E4	4.11E3	0.469
¹⁵⁴ Sm	22.70	82.0	3E-9	E2		1.69E-1	8.88E2	1.5E2	0.542	2.44E4	4.12E3	0.454
⁴⁰ K	0.01	29.8	4.2E-9	(M1)		7.75E-1	1.91E2	1.48E2	0.162	3.59E2	2.78E2	0.139
¹⁶⁸ Er	26.80	79.8	1.9E-9	E2		1.22E-1	1.19E3	1.45E2	0.594	3.25E4	3.98E3	0.498
¹⁷⁴ Hf	0.16	91.0	1.7E-9	E2		1.6E-1	8.94E2	1.43E2	0.632	2.49E4	4E3	0.528
¹⁷⁰ Er	14.90	78.6	1.9E-9	E2		1.16E-1	1.21E3	1.4E2	0.593	3.31E4	3.83E3	0.498
¹⁷⁶ Yb	12.70	82.1	1.8E-9	E2		1.24E-1	1.11E3	1.37E2	0.612	3.04E4	3.76E3	0.513
¹⁸⁷ Os	1.60	74.3	2E-11	M1+E2	0.08	1.18E-1	1.15E3	1.35E2	0.662	1.24E4	1.46E3	0.557
¹⁷² Yb	21.90	78.7	1.6E-9	E2		1.06E-1	1.26E3	1.33E2	0.611	3.43E4	3.64E3	0.513
¹⁶⁰ Gd	21.86	75.3	2.7E-9	E2		1.18E-1	1.09E3	1.28E2	0.557	2.95E4	3.49E3	0.469
¹⁶⁴ Dy	28.20	73.4	2.4E-9	E2		9.93E-2	1.17E3	1.16E2	0.574	3.16E4	3.14E3	0.484
¹⁷⁵ Lu	97.41	113.8	9.9E-11	M1+E2	0.464	2.82E-1	4.07E2	1.15E2	0.630	1.07E4	3.02E3	0.520
¹⁷⁷ Hf	18.61	112.9	5.8E-10	M1+E2	-4.7	3.05E-1	3.76E2	1.15E2	0.639	1.09E4	3.33E3	0.528
¹⁷⁴ Yb	31.80	76.5	1.8E-9	E2		9.5E-2	1.21E3	1.15E2	0.610	3.29E4	3.12E3	0.513
¹⁷⁹ Hf	13.63	122.8	3.7E-11	M1+E2	-0.27	3.06E-1	3.64E2	1.12E2	0.642	7.97E3	2.44E3	0.528
¹⁷³ Yb	16.12	78.6	4.6E-11	M1+E2	-0.224	1.24E-1	8.55E2	1.06E2	0.611	1.88E4	2.33E3	0.513
¹⁰³ Rh	100.00	53.3	1.1E-9	M1		3.21E-1	2.86E2	9.19E1	0.388	5.93E2	1.9E2	0.330
¹⁵³ Eu	52.20	83.4	7.9E-10	M1+E2	0.8	2.07E-1	4.07E2	8.42E1	0.551	1.1E4	2.27E3	0.462
¹⁵³ Eu	52.20	97.4	2E-10	E1		7.62E-1	1.05E2	8.01E1	0.555	2.02E2	1.54E2	0.462

nucleus	nat.ab. %	E_γ , keV	t_e , s	multipole $ e\rangle \rightarrow g\rangle$	δ	f	σ_e , nb 500 keV	σ_γ , nb 500 keV	$Z\alpha E/pc$, 500 keV	σ_e , nb 5 MeV	σ_γ , nb 5 MeV	$Z\alpha E/pc$, 5 MeV
¹⁸³ W	14.30	46.5	1.9E-10	M1+E2	-0.081	1.03E-1	7.57E2	7.8E1	0.637	1.15E4	1.18E3	0.542
¹⁵⁹ Tb	100.00	58.0	5.4E-11	M1+E2	0.119	8.22E-2	9.43E2	7.75E1	0.562	1.67E4	1.37E3	0.476
¹⁶³ Dy	24.90	167.3	3.4E-10	(E2)		4.56E-1	1.67E2	7.59E1	0.605	5.35E3	2.44E3	0.484
¹⁵⁷ Gd	15.65	54.5	1.3E-10	M1+E2	0.19	7.46E-2	9.26E2	6.91E1	0.552	2.06E4	1.54E3	0.469
¹⁵³ Eu	52.20	193.1	2E-10	E2(calc))		4.77E-1	1.34E2	6.38E1	0.589	4.52E3	2.16E3	0.462
⁵⁵ Mn	100.00	126.0	2.6E-10	M1(+E2)	0.052	9.85E-1	5.64E1	5.56E1	0.223	2.75E2	2.71E2	0.183
¹⁵⁵ Gd	14.80	60.0	1.9E-10	M1+E2	-0.197	9.59E-2	5.36E2	5.13E1	0.554	1.19E4	1.14E3	0.469
¹⁶³ Dy	24.90	73.4	1.5E-9	E2		9.95E-2	4.93E2	4.9E1	0.574	1.33E4	1.33E3	0.484
¹⁷¹ Yb	14.30	66.7	8.1E-10	M1+E2	-0.705	7.74E-2	5.95E2	4.6E1	0.607	1.56E4	1.21E3	0.513
¹⁶⁷ Er	22.95	79.3	1.1E-10	M1+E2	-0.2	1.47E-1	3.1E2	4.54E1	0.593	6.5E3	9.52E2	0.498
¹⁸⁷ Os	1.60	187.4	1E-10	E2		1.9E-1	2.29E2	4.35E1	0.707	7.64E3	1.45E3	0.557
¹⁹³ Ir	62.70	139.0	8.2E-11	M1+E2	-0.329	2.95E-1	1.41E2	4.16E1	0.693	3.32E3	9.8E2	0.564
¹⁹¹ Ir	37.30	129.4	1.2E-10	M1+E2	-0.4	2.57E-1	1.52E2	3.89E1	0.690	3.85E3	9.88E2	0.564
¹⁷¹ Yb	14.30	75.9	1.6E-9	E2		6.56E-2	5.69E2	3.73E1	0.610	1.55E4	1.02E3	0.513
¹⁸³ W	14.30	99.1	7.7E-10	E2		8.84E-2	4.1E2	3.62E1	0.652	1.16E4	1.03E3	0.542
¹⁹⁹ Hg	16.87	158.4	2.5E-9	E2		5.22E-1	6.4E1	3.34E1	0.729	2.02E3	1.05E3	0.586
¹⁰⁷ Ag	51.84	32.5	2.9E-9	M1+E2	0.074	7.98E-2	4.18E2	3.33E1	0.402	7.23E3	5.77E2	0.344
⁴⁷ Ti	7.30	159.4	2.1E-10	M1+E2	-0.099	9.94E-1	3.31E1	3.29E1	0.201	2.68E2	2.66E2	0.161
¹⁹ F	100.00	109.9	5.9E-10	E1(calc)		1	3.05E1	3.05E1	0.080	6.13E1	6.13E1	0.066
¹¹⁷ Sn	7.68	158.6	2.8E-10	M1+E2	0.0133	8.63E-1	2.75E1	2.37E1	0.456	8.42E1	7.27E1	0.367
⁶¹ Ni	1.14	67.4	5.3E-9	M1+E2	0.0076	8.9E-1	2.26E1	2.01E1	0.243	5.23E1	4.66E1	0.205
¹⁶⁹ Tm	100.00	118.2	6.2E-11	E2		2.87E-2	6.56E2	1.88E1	0.614	1.92E4	5.51E2	0.506
¹⁸⁹ Os	16.10	69.5	1.6E-9	M1+E2	0.69	8.67E-2	1.97E2	1.71E1	0.660	5.18E3	4.49E2	0.557
¹⁵⁵ Gd	14.80	146.1	1E-10	E2		7.18E-2	2.19E2	1.57E1	0.579	6.76E3	4.85E2	0.469
¹⁵⁷ Gd	15.65	131.4	9.5E-11	E2		4.94E-2	2.81E2	1.39E1	0.574	8.42E3	4.16E2	0.469
¹⁰¹ Ru	17.00	127.2	6.6E-10	M1+E2	0.148	8.55E-1	1.44E1	1.23E1	0.394	2.01E2	1.71E2	0.323
¹⁴⁷ Sm	15.00	121.2	8E-10	M1+E2	-0.33	4.95E-1	2.38E1	1.18E1	0.553	5.69E2	2.82E2	0.454
¹³¹ Xe	21.20	80.2	4.8E-10	M1+E2	0.0165	4.34E-1	2.29E1	9.95	0.471	6.33E1	2.75E1	0.396
²³⁵ U	0.72	46.2	6E-11	M1+E2	0.14	1.72E-2	5.26E2	9.05	0.791	1.09E4	1.88E2	0.674
¹²⁷ I	100.00	57.6	2E-9	M1+E2	-0.084	2.08E-1	4.09E1	8.51	0.458	5.57E2	1.16E2	0.388
¹⁷³ Yb	16.12	179.4	3.2E-11	E2		5.62E-2	1.5E2	8.44	0.647	4.94E3	2.77E2	0.513
¹¹⁹ Sn	8.59	23.9	1.8E-8	M1(+E2)		1.61E-1	4.96E1	7.96	0.426	9.03E1	1.45E1	0.366
¹⁶¹ Dy	18.90	25.7	2.9E-8	E1		3.01E-1	2.6E1	7.82	0.563	3.72E1	1.12E1	0.484
¹⁶⁷ Er	22.95	178.0	5.5E-11	E2(calc)		7.25E-2	1.06E2	7.66	0.628	3.47E3	2.51E2	0.499
¹⁹⁵ Pt	33.80	98.8	1.7E-10	M1+E2	-0.13	1.22E-1	6.26E1	7.62	0.687	8.93E2	1.09E2	0.572
⁹⁹ Ru	12.70	89.7	2E-8	E2+M1	-1.56	4E-1	1.86E1	7.43	0.386	5.14E2	2.06E2	0.323
¹²⁹ Xe	26.40	39.6	9.7E-10	M1+E2	-0.027	7.43E-2	9.62E1	7.15	0.463	5.37E2	3.99E1	0.396
¹⁶⁹ Tm	100.00	8.4	4.1E-9	M1+E2	0.033	5.62E-3	1.19E3	6.71	0.585	2.4E4	1.35E2	0.506
¹⁴⁷ Sm	15.00	197.3	1.2E-9	E2		7.07E-1	8.7	6.16	0.581	2.97E2	2.1E2	0.455
¹²⁵ Te	7.14	35.5	1.5E-9	M1+E2	0.029	6.65E-2	8.57E1	5.7	0.445	5.72E2	3.8E1	0.381
¹⁵⁹ Tb	100.00	137.5	4.1E-11	E2(calc)		2.33E-2	2.39E2	5.58	0.585	7.26E3	1.69E2	0.476
²³² Th	100.00	49.4	3.5E-10	E2		2.97E-3	1.85E3	5.49	0.775	4.8E4	1.43E2	0.660
⁸⁵ Rb	72.17	151.2	7.1E-10	M1+E2	0.072	9.53E-1	5.3	5.05	0.336	3.2E1	3.05E1	0.271
¹⁹⁷ Au	100.00	77.4	1.9E-9	M1+E2	-0.368	1.84E-1	2.71E1	4.99	0.689	6.79E2	1.25E2	0.579
²⁰¹ Hg	13.18	167.4	2.6E-11	M1+E2	0.08	2.43E-1	1.9E1	4.63	0.734	1.22E2	2.97E1	0.586
¹⁵⁵ Gd	14.80	86.5	6.5E-9	E1		6.84E-1	6.6	4.51	0.560	1.22E1	8.36	0.469
²³⁸ U	99.27	44.9	2E-10	E2		1.6E-3	2.73E3	4.36	0.791	7.04E4	1.12E2	0.674
¹²³ Sb	42.64	160.3	6.1E-10	M1+E2	0.063	8.55E-1	4.84	4.14	0.466	2.56E1	2.18E1	0.374

nucleus	nat.ab. %	E_γ , keV	t_e , s	multipole $ e\rangle \rightarrow g\rangle$	δ	f	σ_e , nb 500 keV	σ_γ , nb 500 keV	$Z\alpha E/pc$, 500 keV	σ_e , nb 5 MeV	σ_γ , nb 5 MeV	$Z\alpha E/pc$, 5 MeV
¹⁵⁵ Gd	14.80	105.3	1.2E-9	E1		4.52E-1	8.16	3.69	0.566	1.62E1	7.3	0.469
¹⁴⁵ Nd	8.30	72.5	7.2E-10	M1		2.14E-1	1.56E1	3.35	0.522	3.47E1	7.43	0.440
⁵⁷ Fe	2.20	14.4	9.8E-8	M1+E2	0.00219	1.05E-1	3.05E1	3.19	0.221	5.76E1	6.02	0.191
¹⁸⁷ Os	1.60	75.0	2.2E-9	E2		2.13E-2	1.48E2	3.16	0.662	4.03E3	8.58E1	0.557
⁶⁷ Zn	4.10	184.6	1E-9	M1+E2	0.34	8.51E-1	3.71	3.16	0.279	8.64E1	7.35E1	0.220
⁷⁵ As	100.00	198.6	8.8E-10	M1+E2	0.389	9.81E-1	3.06	3	0.310	7.64E1	7.5E1	0.242
¹⁹³ Ir	62.70	73.0	6.1E-9	M1+E2	-0.558	1.43E-1	1.66E1	2.37	0.670	4.35E2	6.2E1	0.564
¹⁵³ Eu	52.20	151.6	3.6E-10	(E1)		3.35E-1	6.92	2.32	0.572	1.6E1	5.37	0.462
¹⁴¹ Pr	100.00	145.4	1.9E-9	M1+E2	0.069	6.84E-1	3.36	2.3	0.533	1.98E1	1.36E1	0.433
¹⁸⁹ Os	16.10	36.2	5.3E-10	M1+E2	0.045	4.51E-2	4.82E1	2.18	0.651	5.13E2	2.32E1	0.557
¹³³ Cs	100.00	81.0	6.3E-9	M1+E2	-0.151	3.65E-1	5.34	1.95	0.480	9.5E1	3.46E1	0.403
¹⁹⁵ Pt	33.80	199.5	6.6E-10	M1+E2	1.2	1.82E-1	9.78	1.78	0.733	3.22E2	5.87E1	0.572
¹²¹ Sb	57.36	37.1	3.5E-9	M1		8.26E-2	2.09E1	1.72	0.437	4.05E1	3.34	0.374
¹⁴⁹ Sm	13.80	22.5	7.1E-9	M1+E2	0.0715	3.29E-2	5.21E1	1.71	0.529	1.03E3	3.37E1	0.454
⁸³ Kr	11.50	9.4	1.5E-7	M1+E2	0.013	6E-2	2.85E1	1.71	0.305	2.8E2	1.68E1	0.264
¹⁶¹ Dy	18.90	103.1	6E-10	E1		1.78E-1	8.93	1.59	0.583	1.75E1	3.12	0.484
²⁰¹ Hg	13.18	32.2	1E-10	M1+E2	0.017	1.08E-2	1.25E2	1.35	0.684	5.13E2	5.54	0.586
¹⁴⁵ Nd	8.30	67.2	2.9E-8	E2		9.41E-2	1.41E1	1.33	0.521	3.79E2	3.57E1	0.440
²³⁵ U	0.72	103.0	3.3E-11	E2		5.36E-3	2.37E2	1.27	0.812	6.75E3	3.61E1	0.674
¹⁹¹ Ir	37.30	82.4	4.1E-9	M1+E2	-0.88	8.34E-2	1.41E1	1.17	0.673	3.8E2	3.17E1	0.564
¹³⁹ La	99.91	165.9	1.5E-9	M1		7.89E-1	1.48	1.17	0.522	4.47	3.53	0.418
¹³³ Cs	100.00	160.6	1.9E-10	M1+E2	0.96	7.54E-2	1.39E1	1.05	0.502	4.23E2	3.19E1	0.403
¹⁹ F	100.00	197.1	8.9E-8	E2(calc)		9.99E-1	1.03	1.03	0.084	3.51E1	3.51E1	0.066
¹⁵¹ Eu	47.80	21.5	9.6E-9	M1+E2	0.029	3.37E-2	2.95E1	9.94E-1	0.537	3.13E2	1.06E1	0.462
¹⁹⁵ Pt	33.80	129.7	6.7E-10	E2		2.77E-2	3.59E1	9.94E-1	0.699	1.07E3	2.97E1	0.572
²⁰¹ Hg	13.18	26.3	6.3E-10	M1+E2	0.02	1.27E-2	7.38E1	9.39E-1	0.683	4.28E2	5.44	0.586
⁵⁷ Fe	2.20	136.5	8.7E-9	E2		1.04E-1	7.92	8.22E-1	0.234	2.4E2	2.49E1	0.191
¹⁶¹ Dy	18.90	100.4	8.3E-10	(E2)		1.88E-2	4.19E1	7.89E-1	0.582	1.19E3	2.24E1	0.484
¹⁸⁷ Os	1.60	9.7	2.4E-9	M1(+E2)	0	4.31E-3	1.79E2	7.73E-1	0.645	2.99E2	1.29	0.557
¹⁹¹ Ir	37.30	179.0	3.9E-11	M1+E2	-0.75	3.5E-2	1.84E1	6.43E-1	0.712	5.56E2	1.94E1	0.564
¹⁸⁹ Os	16.10	95.3	2.3E-10	M1+E2	0.3	3.17E-2	2.01E1	6.38E-1	0.668	4.76E2	1.51E1	0.557
⁴⁵ Sc	100.00	12.4	3.2E-7	(M2)		2.34E-3	2.26E2	5.28E-1	0.178	9.18E3	2.15E1	0.154
¹⁹³ Ir	62.70	180.1	5.9E-11	M1+E2	-0.48	4.17E-2	8.24	3.44E-1	0.713	2.24E2	9.35	0.564
¹⁶¹ Dy	18.90	74.6	3.1E-9	E1		1.17E-1	1.74	2.03E-1	0.575	3.08	3.59E-1	0.484
¹⁸¹ Ta	99.99	6.2	6E-6	E1		3.23E-2	1.62	5.21E-2	0.619	1.99	6.43E-2	0.535
¹⁵⁷ Gd	15.65	63.9	4.6E-7	E1		2.48E-1	9.63E-2	2.39E-2	0.555	1.64E-1	4.07E-2	0.469
⁶⁷ Zn	4.10	93.3	9.2E-6	E2		5.38E-1	3.18E-2	1.71E-2	0.264	8.91E-1	4.79E-1	0.220
⁷³ Ge	7.73	13.3	3E-6	E2		1.01E-3	6.75	6.81E-3	0.272	1.65E2	1.66E-1	0.235
¹⁵³ Eu	52.20	172.9	1.4E-10	M1+E2	0.77	2.43E-3	4.39E-1	1.07E-3	0.580	1.32E1	3.21E-2	0.462
¹⁵³ Eu	52.20	103.2	3.9E-9	M1+E2	0.127	1.18E-2	7.21E-2	8.49E-4	0.556	9.82E-1	1.16E-2	0.462
¹⁶¹ Dy	18.90	131.8	1.4E-10	E1(calc)		1.37E-3	8.82E-2	1.21E-4	0.592	1.91E-1	2.62E-4	0.484
¹⁷¹ Yb	14.30	122.4	2.7E-7	E2(calc)		1.29E-3	5.73E-3	7.38E-6	0.624	1.69E-1	2.18E-4	0.513
⁷⁷ Se	7.63	161.9	1.7E1	E3		5.29E-1	4.63E-7	2.45E-7	0.311	5.8E-4	3.07E-4	0.249
¹⁵¹ Eu	47.80	196.0	5.9E-5	E3(calc)		1.29E-3	3.88E-5	5E-8	0.590	5.38E-2	6.94E-5	0.462
¹⁰⁹ Ag	48.16	88.0	4E1	E3		3.58E-2	1.26E-6	4.5E-8	0.412	1.27E-3	4.56E-5	0.345
¹⁰⁷ Ag	51.84	93.1	4.4E1	E3		4.64E-2	9.67E-7	4.48E-8	0.413	9.93E-4	4.61E-5	0.345
¹⁰³ Rh	100.00	39.8	3.4E3	E3		6.87E-4	8.72E-8	5.99E-11	0.386	7.71E-5	5.29E-8	0.330
⁸³ Kr	11.50	32.1	6.6E3	E3		4.97E-4	7.31E-9	3.63E-12	0.308	6.33E-6	3.15E-9	0.264